

Wave excitation in the lunar wake associated with solar-wind proton entry

Masaki N. Nishino¹, Yoshifumi Saito¹, Yoshiya Kasahara², Yoshiharu Omura³, Kozo Hashimoto³, Takayuki Ono⁴, Hideo Tsunakawa⁵, Futoshi Takahashi⁵, and Masaki Fujimoto¹

¹ Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo-ku, Sagamihara, 252-5210 Japan, nishino@stp.isas.jaxa.jp

² Kanazawa University, Kakuma-machi, Kanazawa, Ishikawa 920-1192, Japan, kasahara@is.t.kanazawa-u.ac.jp

³ Research Institute for Sustainable Humanosphere, Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan, omura@rish.kyoto-u.ac.jp

⁴ Tohoku University, Aoba, Sendai, Miyagi 980-8578, Japan, ono@stpp.geophys.tohoku.ac.jp

⁵ Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan, htsuna@geo.titech.ac.jp

Abstract

The near-Moon space environment is characterized by formation of a plasma cavity region on the night side along the solar wind (SW) flow, which is called the lunar wake. Because the Moon obstructs the SW plasma flow, the wake region is much more tenuous than the ambient SW region. The plasma environment of the lunar wake has been considered as follows; Because high energy SW electrons can easily access the night side region while SW protons cannot, the wake region should be negatively charged. The electron-rich status of the lunar wake yields ambipolar (inward) electric field at the wake boundary where SW protons gradually come to the wake center along the interplanetary magnetic field (IMF) in the distant wake.

Meanwhile, it has been reported that the SW entry into the distant wake gives rise to plasma waves. Broadband electrostatic noise (BEN) was detected by the Wind spacecraft in the distant wake ($\sim 10,000$ km from the Moon). The excitation of the BEN was attributed to ion two-stream instability owing to the SW proton beams that were accelerated/decelerated into the wake by the ambipolar electric fields. It was also attributed to electron two-stream instability that is expected to occur around the wake boundary region. However, such instabilities were not anticipated to occur in the near-Moon wake (say, ~ 100 km altitude), because it has been believed that a major part of the SW particles cannot access the deepest wake so that any plasma effects would be inoperative there.

Recently, a Japanese lunar orbiter SELENE (Kaguya) performed comprehensive measurements of the plasma and electromagnetic environment around the Moon at ~ 100 km altitude. Concerning the night-side region, two mechanisms of SW proton entry into the near-Moon wake were found. One of them, the type-II entry, which becomes evident when the non-radial IMF component is dominant, is closely related to 'self-pickup' process of protons that were once scattered on the lunar dayside surface or magnetically reflected around the magnetic anomalies. This mechanism lets protons access the deepest region of the wake, namely, $150 < \text{SZA} < 180$ deg at the 100 km altitude of the SELENE orbit.

We found that typical type-II proton entry is accompanied by an enhancement of electron flux and BEN signals. The electron distribution was counter-streaming along the magnetic field, and their energy range was from several hundred eV to 1 keV. This energy is much higher than the typical energy of SW thermal electrons, which shows that acceleration operates as they are attracted from the SW to maintain the quasi-neutrality. Both ends of the magnetic field at the SELENE location were connected to the SW region, which is advantageous to the absorption of SW electrons from both ends into the PGR. The counter-streaming electrons give rise to two-stream instability that results in the excitation of electrostatic solitary waves (ESWs) in the PGR. We also estimate the positive superabundance in the PGR of the deepest wake that yields outward electric field to attract the ambient electrons. The obtained number density ($10^{-5} \sim 10^{-7} \text{ cm}^{-3}$) is orders-of-magnitude lower than the proton density in the PGR ($\sim 0.05 \text{ cm}^{-3}$). This means that the quasi-neutrality is well maintained in the PGR, and that such a tiny superabundance of positive charges can give rise to the prominent feature seen in the electron and plasma wave data. We show that such a coupling of dynamics initiated by the type-II entry leads to excitation of BEN at the bottom of the lunar wake.