



Power Aspects of Processes at the Piston Shock Region

P.A. Sedykh

Irkutsk National Research Technical University, 83, Lermontov str., 664074, Irkutsk, Russian Federation;
e-mail: pvlsd@mail.ru

Abstract

Identifying blast-wave shocks, which can arise during CME formation, is a much more complex problem. The difference from piston shocks is that a blast-wave shock originates from the explosions that frequently accompany CME formation, and further propagates freely without any CME piston effects. Earth's bow shock (BS) is a piston shock. Behind the bow shock front there is a flow of the modified solar wind plasma: transition layer, which also carries the modified magnetic field of solar wind. Velocity and density of plasma as well as parameters of magnetic field of this current can be estimated if the form of the bow shock front and of magnetopause are considered to be known. In this paper we assumed them to be paraboloids of rotation. In this paper we have determined potential distribution along magnetopause from the balance condition of substance coming into transition layer from the solar wind on one side and leaving through the gap between magnetosphere and the bow shock front and through magnetopause on another. To a first approximation this distribution differs from potential distribution at the BS front only in a constant multiplier. We used the established potential distribution as a boundary condition while solving the problem on potential distribution in the magnetosphere. The first solution harmonic turned out to coincide in the form with the boundary condition on magnetopause. We have constructed the full solution for electric field in the magnetosphere having limited ourselves to the first harmonic and corotation field. We have obtained necessary equations to model processes in the region of bow shock.

1 Introduction

Scientific and technical research into shocks, forming in outer space and on the Earth, may be very useful. Shock-wave research was started as far back as the middle of the XIXth century by a group of mathematicians dealing with applied problems and by a group of experimenters. Over the period between the Second World War and until the end of the 20th century, there arose a need for new and accurate data on blast waves from powerful explosions and shocks generated by spacecraft entering dense atmospheric layers and by supersonic flights. The need gave a new impetus for many studies by famous scientists. Vast literature which contains a considerable amount of information, collected in this field of science

for many decades, has been published. A blast-wave shock occurs when in a substance (plasma, gas, liquid, and solid) energy is suddenly released or injected into it from outside, triggering an explosion. For simple explanation, we will consider a spherical explosion. During a spherical explosion, the process proceeds as follows. When energy is suddenly released in the air, water, underground or in plasma, a spherical volume of pressurized hot gas is formed which, when expanding, causes a shock in the environment. The shock moves like a sea tidal wave and immediately raises pressure, density, and temperature of the environment. It also produces a substance flow that follows the shock. Covering ever larger amounts of substance, the shock gradually decays into a weak disturbance or a sound wave. The decay occurs very quickly nearby the explosion center when the shock is very strong and pressure, temperature, density, and velocity behind it are extremely high, and almost ceases when the shock turns into a sound pulse. The sphere, heated by the explosion, is expanding and pulsing until it reaches an equilibrium size corresponding to the environmental pressure. Then the heated substance in the environment diffuses.

Our planet can be represented as a piston moving through the solar wind. Only in this case, we deal with plasma; at the same time we should account for the presence of the interaction between the Earth's magnetic field and the interplanetary magnetic field (frozen-in to the solar wind plasma). The processes occurring in the bow shock region will be even more complex. Earth's bow shock (BS) is a piston shock.

Features of blast-wave and piston shocks are in detail stated in [4]. A method for detecting and identifying piston shocks preceding coronal mass ejections (CMEs) and propagating in various directions relative to the ejections is presented in [4]. The physics behind this method is based on the concept of a perturbed zone [4] that arises in front of the CME, due to its interaction with the background solar wind. When the CME velocity relative to the velocity of the unperturbed solar wind exceeds the local magnetoacoustic speed, which is approximately equal to the Alfvén speed in the solar corona, a piston shock forms in the leading portion of the perturbed zone. In this case, the CME is the piston. Identifying blast-wave shocks, which can also arise during CME formation, is a much more complex problem. The difference from piston shocks is that a blast-wave shock originates from the explosions that frequently

accompany CME formation, and further propagates freely without any CME piston effects.

Since early 1960s many experimental and theoretical investigations of the supersonic and super-Alfvenic solar wind flow past the Earth considered the highly dynamic bow shock boundary upstream of the planetary magnetopause [1]; [11]; [5]; [3]; [2]. A relevant review of pertinent studies can be found in the paper by [7] presenting the empirical model of the Earth's bow shock.

The purpose of this paper is to determine necessary equations for modeling processes in the region of piston shocks. Partially I will use the results from [8-10].

2 Basic equations

Let origins of all coordinate systems (certainly, except for the local) coincide with the origin of the solar-magnetospheric one and let the angle between the tangent to the BS front and axis X be α . Then, we can write the components of interplanetary magnetic field in the local coordinate system as [6], [12]:

$$B_{l0} = B_0[b_{x0}\sin(\varphi/2) + (b_{y0}\sin\psi - b_{z0}\cos\psi)\cos(\varphi/2)] \quad (1)$$

$$B_{n0} = B_0[b_{x0}\cos(\varphi/2) - (b_{y0}\sin\psi - b_{z0}\cos\psi)\sin(\varphi/2)] \quad (2)$$

$$B_{\tau0} = B_0[b_{y0}\cos\psi + b_{z0}\sin\psi]. \quad (3)$$

And current density under the arch of the BS front will be:

$$j_{ll} = -c(\sigma-1)B_{\tau0}/4\pi d \quad (4)$$

$$j_{\tau l} = c(\sigma-1)B_{l0}/4\pi d \quad (5)$$

where d is thickness of the BS front; σ is the ratio between magnetic field tangential components in front of and behind the BS front.




Figure 1. A sketch of the Bow Shock (BS), of the Transition Layer (TL) or magnetosheath, magnetopause (MP). The location of the orthogonal coordinate system (axis x is directed towards the sun, axis y is directed from dawn to dusk and axis z is directed towards the geographic north) and the location of auxiliary coordinate system (r, φ, ψ) and the local coordinate system (l, n, τ).

The corresponding surface current density will be:

$$J_{ll} = -c(\sigma-1)B_{\tau0}/4\pi \quad (6)$$

$$J_{\tau l} = c(\sigma-1)B_{l0}/4\pi \quad (7)$$

These currents' density varies along the front and, therefore, these currents can be divergent. Then there appears a normal current component:

$$j_{nl} = (1/y_g)[\partial(J_{ll}y)/\partial l + \partial J_{\tau l}/\partial\psi] \quad (8)$$

where $y_g = r_g \cdot \cos\varphi$, y is coordinate of the BS front point (r_g, φ) in the auxiliary coordinate system. Index g denotes belonging of the given value to the BS front. In this system the equation of the front looks especially simple:

$$r = y_g/2\cos^2(\varphi/2) \quad (9)$$

where y_g is the distance from the origin of coordinates to the front in the plane YZ, which is equal to the double distance to nose point for the parabola $2x_g$.

In the auxiliary coordinate system the expression for a normal current component will become:

$$j_{nl} = -c[B_{\tau0}\cos^3(\varphi/2)\partial\sigma/\partial\varphi]/2\pi y_g \quad (10)$$

Destiny of this current is subject to examination. In principle, any current in a stationary system must be closed. Only two ways of this current closing can be seen: through magnetosphere body or inside the transition layer. Most likely, they both exist simultaneously. If the current is not used in magnetosphere, it closes in the transition layer, additionally accelerating plasma there. Anyway, this current is ready to work on plasma. Let us notice, by the way, that Poynting vector S is closely connected with electric current density by the ratio:

$$S = j\Phi - (c/4\pi) \operatorname{curl}(B\Phi) \quad (11)$$

Φ – an electric potential.

Now let's turn to determination of electric field at the BS front. Tangential component of this field is:

$$E_{ll} \cong E_{l0} = -V_{n0}B_{\tau0}/c = -(V_0B_{\tau0}/c) \cdot \sin\alpha \quad (12)$$

Integrating E_l along the front on coordinate l we shall obtain:

$$\Psi_g = \Psi_{g0} \cdot \operatorname{tg}(\varphi/2) \quad (13)$$

where $\Psi_{g0} = (V_0B_{\tau0}/c) \cdot y_g$.

As one would expect, the electric field during its movement along the front ($\varphi \rightarrow \pi$) tends to its undisturbed value, with potential tending to infinity. However, the

front itself does not exist far from the disturbing body. Therefore, potential divergence at $\varphi \rightarrow \pi$ should not be taken into account. In reality, we shall simply restrict ourselves to the critical angle of 135° , outside of which we shall consider BS not existing. Now let us estimate magnetopause potential. We shall proceed from the following reasons. After solar wind plasma has crossed the bow shock front, it divides into two flows. One is enclosed in the gap between the BS front and magnetosphere, the second crosses magnetopause and sinks into magnetosphere. It should be noted, that there existed an opinion among geophysicists, that magnetopause is impermeable for plasma. It is easy to show, that the density of electromagnetic energy flow is proportional to the mass flow density. Indeed, having substituted the expression for E into the expression for S , we receive $S = 2Vp_B$, where p_B is magnetic pressure. On the other hand, mass flow density $I = pV$, hence $S = 2I(p_B/\rho)$. Now we can write down the equation of plasma balance:

$$\int n_0 V_{n0} d\Sigma_g = \int n_1 V_{n2} d\Sigma_m + \int n_1 V_{11} d\Sigma_b \quad (14)$$

Here:

$$d\Sigma_g = 2\pi y_g \cdot \text{tg}(\varphi/2) dl_g \quad (15)$$

$$d\Sigma_m = 2\pi y_m \cdot \text{tg}(\varphi/2) dl_m \quad (16)$$

$$d\Sigma_b = \pi(y_g + y_m) \cdot \text{tg}(\varphi/2) dh \quad (17)$$

Where, $d\Sigma$ are elements of the BS front surface, magnetopause and gap area, orthogonal to the flow. We have denoted the width element of the gap through dh .

Correspondingly:

$$n_0 V_{n0} = (cn_0 B_{t0}/B_0^2) \cdot \partial \Psi_g / \partial l_g, \quad (18)$$

$$n_1 V_{n2} = (\sigma cn_0 / B_0) \cdot \partial \Psi_m / \partial l_m, \quad (19)$$

$$n_1 V_{11} = -(cn_0 B_{t0}/B_0^2) \cdot [\partial \Psi_g / \partial h - \partial \Psi_m / \partial h] \quad (20)$$

Substituting (15) - (17) and (18) - (20) into (14) we shall obtain the equation connecting Ψ_g and Ψ_m :

$$\Psi_m = a \int [(\partial \Psi_g / \partial \varphi) \cdot d\varphi] / b \quad (21)$$

where $a = [1 + 2y_g/(y_g + y_m)]$, $b = [1 + 2y_m \sigma B_0^2/(y_g + y_m) B_{t0} B_0]$. Here $B_0^2 = B_{t0}^2 + B_{r0}^2$, and B_0 is magnetic field on magnetopause. In reality b varies within small limits and it can be replaced by average value. Then we can obtain:

$$\Psi_m = (a/b) \cdot \Psi_g \quad (22)$$

Now we can use $\Psi_m(\varphi)$ as a boundary condition for determination of potential inside the magnetosphere.

Since, according to our agreement, magnetopause has the form of a paraboloid of rotation, and magnetic flux tubes are equipotential, then as it was mentioned above, it is enough to receive potential distribution in the equatorial plane. Let us introduce the parabolic coordinate system s, t with the centre in the coordinate origin of the solar-ecliptic one. The equation for potential can be written down in a simple form:

$$\partial^2 \Psi / \partial s^2 + \partial^2 \Psi / \partial t^2 = 0 \quad (23)$$

Any analytical function of the complex number $z = s + it$ will be the solution (23).

Let's represent the solution as power series:

$$\Psi = \sum [A_n (kz^2)^n + B_n (kz^2)^{-n}] \quad (24)$$

We can go over to the Cartesian coordinate system using formulas:

$$x = st, y = (t^2 - s^2)/2$$

It is convenient to present the argument kz^2 in the trigonometrical form: $kz^2 = kr \cdot \exp(i(\varphi + \pi/2))$.

Using function orthogonality, we can find expressions for coefficients:

$$A_n = (1/\pi) \int \Psi_m^0 \cdot \text{tg}(\varphi/2) \cdot (kr)^{-n} \cdot \exp(-i[n(\varphi + \pi/2)]) d\varphi \quad (25)$$

$$B_n = (1/\pi) \int \Psi_m^0 \cdot \text{tg}(\varphi/2) \cdot (kr)^n \cdot \exp(i[n(\varphi + \pi/2)]) d\varphi \quad (26)$$

Taking into account, that $r = y_m/2\cos^2(\varphi/2)$ on magnetopause, we can integrate these expressions (from 0 to π). Since integrals (26) diverge, let us restrict ourselves to determination of real U and imaginary I parts of coefficient A . Further we shall restrict ourselves only to the first decomposition harmonic as it is usually the case while analyzing magnetospheric electric field:

$$\begin{aligned} \Psi_1 &= (2/\pi) \Psi_m^0 (r/y_m) U_1 \sin \varphi \\ &= (2/\pi) \Psi_m^0 (y/y_m) U_1 \end{aligned} \quad (27)$$

It follows from (27), that equipotentials can be lines $y = \text{const}$, that is, lines parallel to axis X . Since magnetospheric electric field is formed by convection and corotation fields, corotation potential should be added to (27):

$$\Psi_c = -\Omega_l R_0^3 B_0 / cr \quad (28)$$

here Ω_l is circular frequency of the Earth rotation, R_0 is radius of the Earth, B_0 is intensity of geomagnetic field at the pole. So, total potential is:

$$\Psi = (2/\pi)\Psi_m^0(r/y_m)U_1 \sin\varphi - \Omega_l R_0^3 B_0/c r. \quad (29)$$

3 Conclusion

It is necessary to note, that interest to research processes in interplanetary shocks and in the bow shock has strongly increased recently (researches within the framework of projects GEOTAIL, CLUSTER-II, THEMIS, SPECTR-R). Selecting and applying of correct initial system of equations are very important for modelling processes in the region of piston shocks. In conclusion, let us remind once again, that we have obtained the necessary equations to model processes in the region of bow shock.

References

1. Beard, D.B. The interaction of the terrestrial magnetic field with the solar corpuscular radiation, *J. Geophys. Res.*, 1960, 65, 3559-3568.
2. Brecht, S.H. Hybrid simulations of the magnetic topology of Mars , *J. Geophys. Res.*, 1997, 102, 4743-4750.
3. Cairns, I.H. and J.G. Lyon. 1995 MHD simulations of the Earth's bow shock at low Mach numbers: Stand-off distances, *J. Geophys. Res.*, 1995, 100, 17173-17182.
4. Eselevich, V.G., M.V. Eselevich, I.V. Zimovets. Blast-wave and Piston Shocks Connected with the Formation and Propagation of a Coronal Mass Ejection. *Astronomy Reports*, 2013. Vol. 57, No. 2, pp. 141 -150. Pleiades Publishing, Ltd..
5. Fairfield, D.H. Average and unusual locations of the Earth's magnetopause and bow shock, *J. Geophys. Res.*, 1971, 76(28), 6700-6716.
6. Korobeinikov V.P. Problems in the Theory of Point Explosion. Moscow. Nauka. Russia. 1985, P.400. (in Russian).
7. Peredo, M., J.A. Slavin, E. Mazur, and S.A. Curtis. Three-dimensional position and Shape of the bow shock and their variation with Alfvénic, sonic and magnetosonic Mach numbers and interplanetary magnetic field orientation, *J. Geophys. Res.*, 1995, 100(A5), 7907-7916.
8. Sedykh P.A. Transformation of solar wind energy into the energy of magnetospheric processes. *Acta Geodaetica et Geophysica*. Springer; 2014. 49, N1, P.1-15. doi: 10.1007/s40328-013-0036-2.
9. Sedykh P.A. Bow shock: Power aspects. *Advances in Space Research*, Elsevier Science. 2014. JASR11746. doi:10.1016/j.asr.2014.03.015.
10. Sedykh P.A. Bow shock: Power aspects. Nova Science Publishers, Inc. In *Horizons in World Physics ed. by Albert Reimer*. NY 11788 USA. 2015. P.53-73.
11. Spreiter, J.R., A.L. Summers, and A.Y. Alksne. Hydromagnetic flow around the magnetosphere, *Planet. Space Sci.*, 1966, 24, 223-253.
12. Whang Y.C. Slow shocks and their transition to fast shocks in the inner solar wind. *J. of Geophys. Res.* 1987, Vol. 92, №5. P.4349-4356.