


## Viability of nowcasting solar flare-driven radio-blackouts using SuperDARN HF radars

Shibaji Chakraborty<sup>\*(1)</sup>, J. Michael Ruohoniemi<sup>(1)</sup> and Joseph B. H. Baker<sup>(1)</sup>

(1) Bradley Department of Electrical and Computer Engineering, Virginia Tech, Blacksburg, USA

The first space weather impact of a solar flare is radio blackout across the dayside of the Earth. At a delay of just 8 minutes, the arrival of enhanced X-ray and EUV radiation leads to a dramatic increase in ionization density in the lower ionosphere [1, 2]. Operation of HF systems are often completely suppressed due to anomalous absorption, while many RF systems suffer some degradation. While the onset of blackout is very rapid ( $\sim 1$ -minute), the recovery takes tens of minutes to hours [3]. Furthermore, severe solar flares can disrupt emergency HF communications that support humanitarian aid services, including amateur radio and satellite communication systems. Our current monitoring capability is based on modeling the ionospheric impacts based on GOES satellite observations of solar fluxes. We present a technique to characterize radio blackout following solar flares using HF radar. The future extension of this work is to develop an now-casting system to identify and monitor radio blackouts using HF radars currently deployed to support space science research. Networks of such radars operate continuously in the northern and southern hemisphere as part of the SuperDARN collaboration. Recent studies have shown that radio blackout (also known as shortwave fadeout) is easily detected and characterized using radar observations. We will combine real-time observations from the North American suite of SuperDARN radars to specify the occurrence of radio blackouts in near real-time.



**Figure 1.** Solar flare-driven SWF monitoring system using SuperDARN HF radar sub-network distributed across the North American sector. Radars near-range fields-of-views are color coded by different phases of SWF given in the figure legend. Yellow color represents no-data.

## References

- [1] D. Watanabe and N. Nishitani, "Study of Ionospheric Disturbances During Solar Flare Events Using the SuperDARN Hokkaido Radar," *Advances in Polar Science*, vol. 24, no. 1, pp. 12–18, 2013.
- [2] S. Chakraborty, J. M. Ruohoniemi, J. B. H. Baker, and N. Nishitani, "Characterization of short-wave fadeout seen in daytime superdarn ground scatter observations," *Radio Science*, vol. 53, no. 4, pp. 472–484, 2018.
- [3] S. Chakraborty, J. B. H. Baker, and J. M. Ruohoniemi, "Probabilistic short-wave fadeout detection in superdarn time series observations," in *2021 IEEE International Conference on Wireless for Space and Extreme Environments (WiSEE)*, 2021, pp. 43–48.