

## **An investigation of whistler intensities above thunderstorms**

*Jiri Fiser*<sup>1</sup>, *Jaroslav Chum*<sup>1</sup>, *Gerhard Diendorfer*<sup>2</sup>, *Michel Parrot*<sup>3</sup>, and *Ondrej Santolik*<sup>4</sup>

<sup>1</sup> Institute Of Atmospheric Physics, Academy of Sciences of the Czech Republic, Bocni II/1401, 14131 Prague, Czech Republic, jifi@ufa.cas.cz

<sup>2</sup> Austrian Electrotechnical Association (OVE-ALDIS), Kahlenberger Str. 2A, 1190 Vienna, Austria, G.Diendorfer@ove.at

<sup>3</sup> LPC2E/CNRS, 3A Avenue de la Recherche Scientifique, 45071 Orléans, France, Michel.Parrot@cnrs-orleans.fr

<sup>4</sup> Charles University, Faculty of Mathematics and Physics, V Holesovickach 2, 18000 Prague, Czech Republic, ondrej.santolik@mff.cuni.cz

### **Abstract**

We study a penetration of whistlers to the ionosphere. We developed an automatic method for assigning causative lightning to fractional-hop whistlers observed on the DEMETER satellite. Processing data from 364 passes of the satellite over Europe, we found that at nighttime, a mean whistler intensity is approximately three times larger than at daytime. A maximum of whistler intensity is shifted approximately one degree from the satellite magnetic footprint owing to the oblique propagation. Calculations of wave attenuation made using IRI2007 and MSIS models show very similar result in a ratio of nighttime and daytime intensities.